

K-THEORY FOR CUNTZ-KRIEGER ALGEBRAS ARISING FROM REAL QUADRATIC MAPS

NUNO MARTINS, RICARDO SEVERINO, AND J. SOUSA RAMOS

ABSTRACT. We compute the K -groups for the Cuntz-Krieger algebras $\mathcal{O}_{A_{\mathcal{K}(f_\mu)}}$, where $A_{\mathcal{K}(f_\mu)}$ is the Markov transition matrix arising from the *kneading sequence* $\mathcal{K}(f_\mu)$ of the one-parameter family of real quadratic maps f_μ .

Consider the one-parameter family of real quadratic maps $f_\mu : [0, 1] \rightarrow [0, 1]$ defined by $f_\mu(x) = \mu x(1 - x)$, with $\mu \in [0, 4]$. Using Milnor-Thurston's kneading theory [14], J. Guckenheimer [5] has classified up to topological conjugacy, a certain class of maps which includes the quadratic family. The idea of kneading theory is to encode information about the orbits of a map in terms of infinite sequences of symbols and to exploit the natural order of the interval to establish topological properties of the map. In the following, I will denote the unit interval $[0, 1]$ and c the unique turning point of f_μ . For $x \in I$, let

$$\varepsilon_n(x) = \begin{cases} -1 & \text{if } f_\mu^n(x) > c, \\ 0 & \text{if } f_\mu^n(x) = c, \\ +1 & \text{if } f_\mu^n(x) < c. \end{cases}$$

The sequence $\varepsilon(x) = (\varepsilon_n(x))_{n=0}^\infty$ is called the *itinerary* of x . The itinerary of $f_\mu(c)$ is called the *kneading sequence* of f_μ and will be denoted by $\mathcal{K}(f_\mu)$. Observe that $\varepsilon_n(f_\mu(x)) = \varepsilon_{n+1}(x)$, i.e. $\varepsilon(f_\mu(x)) = \sigma\varepsilon(x)$ where σ is the shift map. Let $\Sigma = \{-1, 0, +1\}$ be the alphabet set. The sequences on $\Sigma^\mathbb{N}$ are ordered lexicographically. However, this ordering is not reflected by the mapping $x \rightarrow \varepsilon(x)$, because the map f_μ reverses orientation on $[c, 1]$. To take this into account, for a sequence $\varepsilon = (\varepsilon_n)_{n=0}^\infty$ of the symbols $-1, 0, +1$, another sequence $\theta = (\theta_n)_{n=0}^\infty$ is defined by $\theta_n = \prod_{i=0}^n \varepsilon_i$. If $\varepsilon = \varepsilon(x)$ is the itinerary of a point $x \in I$ then $\theta = \theta(x)$ is called the *invariant coordinate* of

Date: September 19, 2002.

1991 Mathematics Subject Classification. Primary 37A55, 37B10, 37E05; Secondary 46L80.

Key words and phrases. Cuntz-Krieger algebras, real quadratic maps, K -groups.

x . The fundamental observation of J. Milnor and W. Thurston [14] is the monotonicity of the invariant coordinates:

$$x < y \Rightarrow \theta(x) \leq \theta(y).$$

Let us now consider only those kneading sequences that are periodic, i.e.

$$\begin{aligned} \mathcal{K}(f_\mu) &= \varepsilon_0(f_\mu(c)) \dots \varepsilon_{n-1}(f_\mu(c)) \varepsilon_0(f_\mu(c)) \dots \varepsilon_{n-1}(f_\mu(c)) \dots \\ &= (\varepsilon_0(f_\mu(c)) \dots \varepsilon_{n-1}(f_\mu(c)))^\infty \equiv (\varepsilon_1(c) \dots \varepsilon_n(c))^\infty \end{aligned}$$

for some $n \in \mathbb{N}$. The sequences $\sigma^i(\mathcal{K}(f_\mu)) = \varepsilon_{i+1}(c) \varepsilon_{i+2}(c) \dots$, $i = 0, 1, 2, \dots$, will then determine a Markov partition of I into $n-1$ line intervals $\{I_1, I_2, \dots, I_{n-1}\}$ [15], whose definitions will be given in the proof of Theorem 1. Thus, we will have a Markov transition matrix $A_{\mathcal{K}(f_\mu)}$ defined by

$$A_{\mathcal{K}(f_\mu)} := (a_{ij}) \text{ with } a_{ij} = \begin{cases} 1 & \text{if } f_\mu(\text{int } I_i) \supseteq \text{int } I_j \\ 0 & \text{otherwise.} \end{cases}$$

It is easy to see that this matrix $A_{\mathcal{K}(f_\mu)}$ is not a permutation matrix and no row or column of $A_{\mathcal{K}(f_\mu)}$ is zero. Thus, for each one of these matrices and following the work of J. Cuntz and W. Krieger [2], one can construct the Cuntz-Krieger algebra $\mathcal{O}_{A_{\mathcal{K}(f_\mu)}}$. In [3], J. Cuntz proved that

$$K_0(\mathcal{O}_A) \cong \mathbb{Z}^r / (1 - A^T) \mathbb{Z}^r \quad \text{and} \quad K_1(\mathcal{O}_A) \cong \ker(I - A^t : \mathbb{Z}^r \rightarrow \mathbb{Z}^r).$$

for a $r \times r$ matrix A that satisfies a certain condition (I) (see [2]), which is readily verified by the matrices $A_{\mathcal{K}(f_\mu)}$. In [1] R. Bowen and J. Franks introduced the group $BF(A) := \mathbb{Z}^r / (1 - A) \mathbb{Z}^r$ as an invariant for flow equivalence of topological Markov subshifts determined by A .

We can now state and prove the following.

Theorem 1. *Let $\mathcal{K}(f_\mu) = (\varepsilon_1(c) \varepsilon_2(c) \dots \varepsilon_n(c))^\infty$, for some $n \in \mathbb{N} \setminus \{1\}$. Thus, we have*

$$K_0(\mathcal{O}_{A_{\mathcal{K}(f_\mu)}}) \cong \mathbb{Z}_a \quad \text{with} \quad a = \left| 1 + \sum_{l=1}^{n-1} \prod_{i=1}^l \varepsilon_i(c) \right|$$

and

$$K_1(\mathcal{O}_{A_{\mathcal{K}(f_\mu)}}) \cong \begin{cases} \{0\} & \text{if } a \neq 0 \\ \mathbb{Z} & \text{if } a = 0. \end{cases}$$

Proof. Set $z_i = \varepsilon_i(c) \varepsilon_{i+1}(c) \dots$ for $i = 1, 2, \dots$. Let $z'_i = f_\mu^i(c)$ be the point on the unit interval $[0, 1]$ represented by the sequence z_i for $i = 1, 2, \dots$. We have $\sigma(z_i) = z_{i+1}$ for $i = 1, \dots, n-1$ and $\sigma(z_n) = z_1$. Denote by ω the $n \times n$

matrix representing the shift map σ . Let C_0 be the vector space spanned by the formal basis $\{z'_1, \dots, z'_n\}$. Now, let ρ be the permutation of the set $\{1, \dots, n\}$, which allows us to order the points $z'_1, \dots, z'_n$ on the unit interval $[0, 1]$, i.e.

$$0 < z'_{\rho(1)} < z'_{\rho(2)} < \dots < z'_{\rho(n)} < 1.$$

Set $x_i := z'_{\rho(i)}$ with $i = 1, \dots, n$ and let π denote the permutation matrix which takes the formal basis $\{z'_1, \dots, z'_n\}$ to the formal basis $\{x_1, \dots, x_n\}$. We will denote by C_1 the $n - 1$ dimensional vector space spanned by the formal basis $\{x_{i+1} - x_i : i = 1, \dots, n - 1\}$. Set

$$I_i := [x_i, x_{i+1}], \quad \text{for } i = 1, \dots, n - 1.$$

Thus, we can define the Markov transition matrix $A_{\mathcal{K}(f_\mu)}$ as above. Let φ denote the incidence matrix that takes the formal basis $\{x_1, \dots, x_n\}$ of C_0 to the formal basis $\{x_2 - x_1, \dots, x_n - x_{n-1}\}$ of C_1 . Put $\eta := \varphi\pi$. As in [7] and [8], we obtain an endomorphism α of C_1 , that makes the following diagram commutative.

$$\begin{array}{ccccc} & & \eta & & \\ & C_0 & \longrightarrow & C_1 & \\ \omega \downarrow & & & & \downarrow \alpha \\ & C_0 & \longrightarrow & C_1 & \\ & & \eta & & \end{array}$$

We have $\alpha = \eta\omega\eta^T(\eta\eta^T)^{-1}$. Remark that if we neglect the negative signs on the matrix α then we will obtain precisely the Markov transition matrix $A_{\mathcal{K}(f_\mu)}$. In fact, consider the $(n - 1) \times (n - 1)$ matrix

$$\beta := \begin{bmatrix} 1_{n_L} & 0 \\ 0 & -1_{n_R} \end{bmatrix}$$

where 1_{n_L} and 1_{n_R} are the identity matrices of rank n_L and n_R respectively, with n_L (n_R) being the number of intervals I_i of the Markov partition placed on the left (right) hand side of the turning point of f_μ . Therefore, we have

$$A_{\mathcal{K}(f_\mu)} = \beta\alpha.$$

Now, consider the following matrix defined by

$$\gamma_{\mathcal{K}(f_\mu)} := (\gamma_{ij}) \quad \text{with} \quad \begin{cases} \gamma_{ii} = \varepsilon_i(c), & i = 1, \dots, n \\ \gamma_{in} = -\varepsilon_i(c), & i = 1, \dots, n \\ \gamma_{ij} = 0, & \text{otherwise.} \end{cases}$$

The matrix $\gamma_{\mathcal{K}(f_\mu)}$ makes the diagram

$$\begin{array}{ccc} & \eta & \\ & C_0 \longrightarrow C_1 & \\ \gamma_{\mathcal{K}(f_\mu)} \downarrow & & \downarrow \beta \\ & C_0 \longrightarrow C_1 & \\ & \eta & \end{array}$$

commutative. Finally, set $\theta_{\mathcal{K}(f_\mu)} := \gamma_{\mathcal{K}(f_\mu)} \omega$. Then, the following diagram

$$\begin{array}{ccc} & \eta & \\ & C_0 \longrightarrow C_1 & \\ \theta_{\mathcal{K}(f_\mu)} \downarrow & & \downarrow A_{\mathcal{K}(f_\mu)} \\ & C_0 \longrightarrow C_1 & \\ & \eta & \end{array}$$

is also commutative. Now, notice that the transpose of η has the following factorization

$$\eta^T = Y i X,$$

where Y is an invertible (over $\mathbb{Z}$) $n \times n$ integer matrix given by

$$Y := \begin{pmatrix} 1 & 0 & \cdots & & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ \vdots & 0 & \ddots & \ddots & \vdots \\ & \vdots & & & 0 \\ 0 & 0 & \cdots & 0 & 1 & 0 \\ -1 & -1 & \cdots & & -1 & 1 \end{pmatrix},$$

i is the inclusion $C_1 \hookrightarrow C_0$ given by

$$i := \begin{pmatrix} 1 & 0 & & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & & 0 \\ 0 & \cdots & & 1 \\ 0 & \cdots & & 0 \end{pmatrix}$$

and X is an invertible (over $\mathbb{Z}$) $(n-1) \times (n-1)$ integer matrix obtained from the $(n-1) \times n$ matrix η^T by removing the n -th row of η^T . Thus, from the commutative diagram

$$\begin{array}{ccc} & \eta^T & \\ & C_1 \longrightarrow C_0 & \\ A_{\mathcal{K}(f_\mu)}^T \downarrow & & \downarrow \theta_{\mathcal{K}(f_\mu)}^T \\ & C_1 \longrightarrow C_0 & \\ & \eta^T & \end{array}$$

we will have the following commutative diagram with short exact rows

$$\begin{array}{ccccccc}
0 & \longrightarrow & C_1 & \xrightarrow{i} & C_0 & \xrightarrow{p} & C_0/C_1 \longrightarrow 0 \\
& & \downarrow A' & & \downarrow \theta' & & \downarrow 0 \\
0 & \longrightarrow & C_1 & \xrightarrow{i} & C_0 & \xrightarrow{p} & C_0/C_1 \longrightarrow 0
\end{array}$$

where the map p is represented by the $1 \times n$ matrix $\begin{bmatrix} 0 & \dots & 0 & 1 \end{bmatrix}$ and

$$A' = X A_{\mathcal{K}(f_\mu)}^T X^{-1} \quad \text{and} \quad \theta' = Y^{-1} \theta_{\mathcal{K}(f_\mu)}^T Y$$

i.e., A' is similar to $A_{\mathcal{K}(f_\mu)}^T$ over $\mathbb{Z}$ and θ' is similar to $\theta_{\mathcal{K}(f_\mu)}^T$ over $\mathbb{Z}$. Hence, for example by [10], we obtain respectively

$$\begin{aligned}
\mathbb{Z}^{n-1}/(1 - A')\mathbb{Z}^{n-1} &\cong \mathbb{Z}^{n-1}/(1 - A_{\mathcal{K}(f_\mu)})\mathbb{Z}^{n-1} \quad \text{and} \\
\mathbb{Z}^n/(1 - \theta')\mathbb{Z}^n &\cong \mathbb{Z}^n/(1 - \theta_{\mathcal{K}(f_\mu)})\mathbb{Z}^n.
\end{aligned}$$

Now, from the last diagram we have, for example by [9],

$$\theta' = \begin{bmatrix} A' & * \\ 0 & 0 \end{bmatrix}.$$

Therefore,

$$\mathbb{Z}^{n-1}/(1 - A')\mathbb{Z}^{n-1} \cong \mathbb{Z}^n/(1 - \theta')\mathbb{Z}^n$$

and

$$\mathbb{Z}^{n-1}/(1 - A_{\mathcal{K}(f_\mu)})\mathbb{Z}^{n-1} \cong \mathbb{Z}^n/(1 - \theta_{\mathcal{K}(f_\mu)})\mathbb{Z}^n.$$

Next, we will compute $\mathbb{Z}^n/(1 - \theta_{\mathcal{K}(f_\mu)})\mathbb{Z}^n$. From the previous discussions and notations, the $n \times n$ matrix $\theta_{\mathcal{K}(f_\mu)}$ is explicitly given by

$$\theta_{\mathcal{K}(f_\mu)} := \begin{pmatrix} -\varepsilon_1(c) & \varepsilon_1(c) & 0 & \dots & 0 \\ \vdots & 0 & \ddots & \ddots & \vdots \\ & \vdots & \ddots & & 0 \\ -\varepsilon_{n-1}(c) & & & & \varepsilon_{n-1}(c) \\ 0 & 0 & \dots & & 0 \end{pmatrix}.$$

Notice that this matrix $\theta_{\mathcal{K}(f_\mu)}$ completely describes the dynamics of f_μ . Finally, using row and column elementary operations over $\mathbb{Z}$, we can find

invertible (over $\mathbb{Z}$) matrices U_1 and U_2 with integer entries such that

$$1 - \theta_{\mathcal{K}(f_\mu)} = U_1 \begin{pmatrix} 1 + \sum_{l=1}^{n-1} \prod_{i=1}^l \varepsilon_i(c) & & & & \\ & 1 & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & 1 \end{pmatrix} U_2.$$

Thus, we obtain

$$K_0(\mathcal{O}_{A_{\mathcal{K}(f_\mu)}}) \cong \mathbb{Z}^{n-1} / (1 - A_{\mathcal{K}(f_\mu)}^T) \mathbb{Z}^{n-1} \cong \mathbb{Z}_a,$$

where $a = \left| 1 + \sum_{l=1}^{n-1} \prod_{i=1}^l \varepsilon_i(c) \right|$ and $n \in \mathbb{N} \setminus \{1\}$. ■

Example 1. *Set*

$$\mathcal{K}(f_\mu) = (RLLRRC)^\infty,$$

where $R = -1$, $L = +1$, $C = 0$. Thus, we can construct the 5×5 Markov transition matrix $A_{\mathcal{K}(f_\mu)}$ and the matrices $\theta_{\mathcal{K}(f_\mu)}$, ω , φ , and π .

$$A_{\mathcal{K}(f_\mu)} = \begin{pmatrix} 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 & 0 \end{pmatrix}, \quad \theta_{\mathcal{K}(f_\mu)} = \begin{pmatrix} 1 & -1 & 0 & 0 & 0 \\ -1 & 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$\omega = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad \varphi = \begin{pmatrix} -1 & 1 & & & \\ & -1 & 1 & & \\ & & -1 & 1 & \\ & & & -1 & 1 \\ & & & & -1 & 1 \end{pmatrix},$$

$$\pi = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

We have

$$K_0(\mathcal{O}_{A_{\mathcal{K}(f_\mu)}}) \cong \mathbb{Z}_2 \quad \text{and} \quad K_1(\mathcal{O}_{A_{\mathcal{K}(f_\mu)}}) \cong \{0\}.$$

Remark 1. In the statement of Theorem 1 the case $a = 0$ may occur. This happens when we have a star product factorizable kneading sequence [4]. In this case the correspondent Markov transition matrix is reducible.

Remark 2. In [6] the authors have constructed a class of C^* -algebras from the β -expansions of real numbers. In fact, by considering a semiconjugacy from the real quadratic map to the tent map [14], we can also obtain Theorem 1 using [6] and the λ -expansions of real numbers introduced in [4].

Remark 3. In [12] (see also [11]) and [13] the BF-groups are explicitly calculated with respect to other kind of maps on the interval.

REFERENCES

- [1] R. Bowen and J. Franks. Homology for zero dimensional basic sets. *Ann. Math. (2)* **106** (1977), 73-92.
- [2] J. Cuntz and W. Krieger. A class of C^* -algebras and topological Markov chains. *Invent. Math.* **56** (1980), 251-268.
- [3] J. Cuntz. A class of C^* -algebras and topological Markov chains II: Reducible chains and the Ext-functor for C^* -algebras. *Invent. Math.* **63** (1981), 25-40.
- [4] B. Derrida, A. Gervois and Y. Pomeau. Iteration of endomorphisms on the real axis and representations of numbers. *Ann. Inst. H. Poincaré Sect. A (N.S.)* **29** (1978), 305-356.
- [5] J. Guckenheimer. Sensitive Dependence to Initial Conditions for One Dimensional Maps. *Comm. Math. Phys.* **70** (1979), 133-160.
- [6] Y. Katayama, K. Matsumoto and Y. Watatani. Simple C^* -algebras arising from β -expansion of real numbers. *Ergodic Theory Dynam. Systems* **18** (1998), 937-962.
- [7] J. P. Lampreia and J. Sousa Ramos. Trimodal maps. *Internat. J. Bifur. Chaos Appl. Sci. Engrg.* **3** (1993), 1607-1617.
- [8] J. P. Lampreia and J. Sousa Ramos. Symbolic dynamics for bimodal maps. *Portugal. Math.* **54** (1997), 1-18.
- [9] S. Lang. *Algebra*. Addison-Wesley, 1965.
- [10] D. Lind and B. Marcus. *An Introduction to Symbolic Dynamics and Coding*. Cambridge University Press, 1995.
- [11] N. Martins and J. Sousa Ramos. Cuntz-Krieger algebras arising from linear mod one transformations. *Fields Inst. Commun.* **31** (2002), 265-273.
- [12] N. Martins and J. Sousa Ramos. Bowen-Franks groups associated with linear mod one transformations. To appear in *Internat. J. Bifur. Chaos Appl. Sci. Engrg. (2003)*.
- [13] N. Martins, R. Severino and J. Sousa Ramos. Bowen-Franks groups for bimodal matrices. To appear in *J. Differ. Equations Appl.*
- [14] J. Milnor and W. Thurston. On iterated maps of the interval. *Lecture Notes in Math.* **1342** (1988), 465-563.
- [15] P. Stefan. A theorem of Sharkovsky on the existence of periodic orbits of continuous endomorphisms of the real line. *Comm. Math. Phys.* **54** (1977), 237-248.

DEPARTAMENTO DE MATEMÁTICA, INSTITUTO SUPERIOR TÉCNICO

E-mail address: `nmartins@math.ist.utl.pt`

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE DO MINHO

E-mail address: `ricardo@math.uminho.pt`

DEPARTAMENTO DE MATEMÁTICA, INSTITUTO SUPERIOR TÉCNICO

E-mail address: `sramos@math.ist.utl.pt`